

Accepted Manuscript

Smoking status in Iranian male adolescents: A cross-sectional study and a meta-analysis

Milad Nazarzadeh, Zeinab Bidel, Erfan Ayubi, Abolfazl Bahrami, Fateme Jafarie, Ashgar Mohammadpoorasl, Ali Delpisheh, Farhad Taromian

PII: S0306-4603(13)00030-0  
DOI: doi: [10.1016/j.addbeh.2013.01.018](https://doi.org/10.1016/j.addbeh.2013.01.018)  
Reference: AB 3882

To appear in: *Addictive Behaviors*



Please cite this article as: Nazarzadeh, M., Bidel, Z., Ayubi, E., Bahrami, A., Jafarie, F., Mohammadpoorasl, A., Delpisheh, A. & Taromian, F., Smoking status in Iranian male adolescents: A cross-sectional study and a meta-analysis, *Addictive Behaviors* (2013), doi: [10.1016/j.addbeh.2013.01.018](https://doi.org/10.1016/j.addbeh.2013.01.018)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Smoking status in Iranian male adolescents: A cross-sectional study and a meta-analysis

**Milad Nazarzadeh**, MSc of Epidemiology, Psychosocial Injuries Research Center, Ilam University of Medical Sciences, Ilam, Iran . Email: Nazarzadeh\_milad@yahoo.com.

**Zeinab Bidel**, MSc of Epidemiology, Department of Clinical Epidemiology, Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran. Email: Zeinab.bidel@yahoo.com.

**Erfan Ayubi**, MSc of Epidemiology, Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: aubi65@gmail.com.

**Abolfazl Bahrami**, B.S of Public Health, School of Health, Student Research Committee, Zanzan University of Medical Sciences, Iran. Email: abolfazlbahrami30@yahoo.com.

**Fateme Jafari**, Assistant Professor of Maternal and Child Health, School of Health, Department of Public Health, Zanzan University of Medical Sciences, Zanzan, Iran. Email: fjafari@zums.ac.ir.

**Ashgar Mohammadpoorasl**, PhD of Epidemiology, Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Science, Tehran, Iran. Email: poorasl@yahoo.com.

**Ali Delpisheh**, Professor of Clinical Epidemiology, Department of Clinical Epidemiology, Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran. Email: alidelpisheh@yahoo.com.

**Farhad Taromian**, Associate Professor of Clinical Psychology, School of Medicine, Zanzan University of Medical Sciences, Zanzan, Iran. Email: taremian@yahoo.com.

**Corresponding author:** Prof. Ali Delpisheh (PhD, PostDoc)

Department of Clinical Epidemiology  
Faculty of Medicine  
Ilam University of Medical Sciences,  
Ilam, Iran  
Telefax: +98-841-2227103

Email: [alidelpisheh@yahoo.com](mailto:alidelpisheh@yahoo.com)

## ABSTRACT

The present study aimed to estimate the prevalence of smoking status and its associated factors in Iranian adolescents and a meta-analysis of recent cross-sectional studies in order to estimate the corresponding prevalence for all Iranian adolescents.

In a cross-sectional study, 1064 male high school students in Zanjan city (northwest of Iran) were recruited. A self-administered questionnaire was used for smoking status and associated factors. Through the meta-analysis, all relevant published studies were reviewed.

Almost one-third of adolescents (34.2%, n= 354) has experienced smoking either experimentally (23.4%, n=242), or regularly (10.8%, n=112). Multivariate analysis showed that older age (OR=1.20; 95% CI: 1.05-1.37), risky behaviors (OR=1.83; 1.25-2.68), Tramadol medication (OR=2.19; 1.54-3.11), low self-esteem (OR=1.07; 1.03-1.11), positive attitude toward smoking (OR=1.15; 1.09-1.21), positive thinking about smoking (OR=1.07; 1.01-1.14) and having smoker friends (OR=1.94; 1.36-2.77) were significantly associated with cigarette smoking in adolescents. Meta-analysis results showed that 7 % of Iranian adolescents are regular smokers and 27 % are experimenters.

Increasing prevalence of smoking in Iranian adolescents is a major concern for public health. Controlling risky behaviors and increasing health education are recommended.

**Keyword:** Adolescents, Iranian, smoking status, meta-analysis

## 1. INTRODUCTION

Smoking is one of the important causes of public health problems around the world (Delpisheh, Attia, Drammond, & Brabin, 2006). One of the most significant challenges for smoking cessation programs is prevention of smoking onset during mid or late adolescence (Hiemstra, Otten, & Engels, 2012). It has been revealed that distal and proximal risk factors including smoker peers, socioeconomic status (SES), family condition and risk-taking behaviors affect the probability of start smoking in adolescents (Delpisheh, Kelly, Rizwan, & Brabin, 2006) (Wang, Ho, Lo, & Lam, 2012) (Schaefer, Haas, & Bishop, 2012). There are debates about recently emerged risk factor such as Tramadol and other prescription medications for adolescent smoking. Despite, effect of these drugs on abuser people (Epstein, Preston, & Jasinski, 2006) (Farajidana, Hassanian-Moghaddam, Zamani, & Sanaei-Zadeh, 2012) (Marechal, Honorat, & Claudet, 2011), its association with adolescent smoking is of grate concerns for public health.

Twenty five European countries studied, showed a prevalence of 22% for adolescent smoking in males (Baska, Warren, Baskova, & Jones, 2009), which is obviously higher than the corresponding rate for Iranian adolescents from 2.5 to 17% (Alireza Ayatollahi, Mohammadpoorasl, & Rajaeifard, 2005) (Mohammadpoorasl, Fakhari, Shamsipour, Rostami, & Rashidian, 2011) (Kelishadi, et al., 2006). Reviewing cross-sectional studies in Iran showed different prevalence rates across the country. So far, no meta-analysis has been conducted in Iran to assess differences between provinces in terms of adolescent smoking and calculate a combined prevalence.

The present study aimed to calculate adolescent smoking prevalence in Zanjan city (northwest of Iran) in the 2011-2012 school year and to examine the psychological, pharmaceutical and social factors associated with adolescent smoking. A meta-analysis of recent cross-sectional studies in Iran was also conducted in order to estimate adolescent smoking in Iran.

## 2. MATERIALS & METHODS

Study participants were male students who were randomly recruited from 61 classes out of 42 high schools in Zanjan city. Initially, about 12000 high school students at grades 9<sup>th</sup>, 10<sup>th</sup>, 11<sup>th</sup> and 12<sup>th</sup> from 42 high schools were identified. Of them 1100 (9.1%) were selected by cluster and stratified samplings.

The questionnaire design, type of variables, variables scales and its reliability and validity has been reported in detail elsewhere (Alireza Ayatollahi, et al., 2005). Smoking prevalence was determined using a valid algorithm as never smokers, experimenter and regular smokers accordingly (Kaplan, Napoles-Springer, Stewart, & Perez-Stable, 2001).

To detect the prescription medications status, the following direct question was added to the questionnaire; “Have you ever used either Tramadol, or Ritalin?”

The general risk taking behavior was measured according to Kaplan method (Kaplan, et al., 2001) and self-esteem was measured using the Rosenberg 10-item questionnaire (Alireza Ayatollahi, et al., 2005). The attitude toward smoking was measured according to Hill (Hill, Boudreau, Amyot, Dery, & Godin, 1997). To investigate the reaction of relatives and closest friends about smoking behaviors, the following question was asked; “If you lit up a cigarette in front of your father/mother/brother/sister/closest friend, how he/or she would react?” Two severe and mild reactions were considered for the answers. Socio-economic status (SES) was built through relevant characteristics such as parental education, house furniture and family income.

A logistic regression model was used to calculate adjusted odds ratios and chi-square test was used to examine the association between smoking status and independent variables. The one-way analysis of variance (ANOVA) was used to compare mean score of self-esteem, attitude toward smoking and positive thinking about smoking and two-way analysis of variance (MANOVA) was used to assess interested interaction.

For the meta-analysis, all available data concerning adolescent smoking in Iran were investigated. The search was done on May 15, 2012. Studies were found with

electronically searched MEDLINE, EMBASE, and the ISI the Web of Science using the following search strategy: "Adolescents" OR "Teen" OR "Teenager" OR "Youth" AND "Smoking" OR "Cigarette smoking" OR "Tobacco smoking" AND "Iran". Meanwhile, all national databases including Magiran, IranDoc, SID, IranPsych and IranMedex were systematically searched. Finally, data from 27 studies was prepared and included. It was considered prevalence of regular and experimental smoking in adolescents as interested measure.

Heterogeneity was assessed using chi-square test with significance level of  $< 0.1$  and the  $I^2$  statistic for estimation of inconsistency in meta-analyses. The random effect model was used to calculate the pooled prevalence and Egger test for evaluation of publication bias. All analyses were carried out using Stata SE 11.2 and SPSS (version 20).

### 3. RESULTS

#### 3.1. Cross-sectional study:

The response rate was 94% (1035/1100). Mean age was  $17.2 \pm 1.3$  years ranged between 14 to 21 years. Out of 354 (34.2%) adolescent smokers, 242 (23.4%) were classified as experimenters and 112 (10.8%) as regular smokers. Overall, 404 students (42.5%) had at least one friend who smoked. In terms of medications, 73 (36.0%) and 50 (4.8%) students had used either Tramadol and/or Ritalin respectively. Mean self-esteem scores in never smokers, experimenter and regular smokers were  $17.2 \pm 4.9$ ,  $19.4 \pm 4.9$  and  $19.3 \pm 5.4$  respectively ( $p < 0.001$ ). Mean score of positive attitude toward smoking in never smokers, experimenters and regular smokers were  $-10.4 \pm 3.2$ ,  $-7.8 \pm 4.9$ , and  $-4.01 \pm 6.4$  respectively ( $p < 0.001$ ). Majority of students (78.2 %) reported easy access to cigarettes. Demographic characteristics are presented in Table 1.

The logistic regression revealed that older age, having general risk taking behaviors, Tramadol, lower self-esteem, positive attitude toward smoking, positive thinking about smoking and having smoker friend were significantly associated with cigarette smoking in adolescents (Table 2).

### 3.2. Meta-analysis results

We found 27 relevant studies (A Mohammadpoorasl, et al., 2011) (Habib E, Shirzad AS, Naseri-Kouzehgarani G, Hooman S, & Reza MM, 2012) (Alaei Khorayem R, Kadeyor P, Mohamadkhani Sh, Sarami GhR, & Alaei Khorayem S, 2011) (Ramzankhani A, Sarbandi Zaboli F, Zargi A, Heydari GhR, & Masjedi MR, 2010) (Rahmaniyan KA, Jafarzadeh A, & Khaloei A, 2010) (Najafi K, Fekri F, & etal, 2007) (Namakin K, Sharifzadeh Gh, & Miri MR, 2008) (Mohammadpoor Asl, Fakhari, Rostami, & Pourafkary, 2007) (Barikani A, 2008) (Ziaaddini H, Kheradmand A, Nakhaei N, & Taherzadeh H, 2008) (Ziaaddini H, et al., 2008) (Miri M, Moasheri B, & Hanafi H, 2004) (Mohtasham Amiri Z, 2008) (Mojahed A & Bakhshani Nm, 2004) (Heydari, Sharifi, Hosseini, & Masjedi, 2007) (Alireza Ayatollahi, et al., 2005) (Aslinejad M, Alemi A, & Chamanzari H, 2004) (Mokhtari Habashi Gh, Bahrani N, & SobhRakhshankhah M, 2001) (Jafarzadegan T & Zarehei F, 2001) (Ahmadi & Hasani, 2003) (Ahmadi & Ostovan, 2002) (Aghaei A, 2001) (Ziaei P, 2001) (Shariat Zadeh M & Sadeghi A, 2001) (Hamidzadeh Y & Hosieni Ahog MM, 2001) (Najafi, et al., 2004). In 14 studies and in 26 studies enough data were available to calculate the prevalence of regular and experimental smoking respectively. Sample sizes in meta-analysis of experimental smoking were 26012 and 19732 for regular smoking.

There were evidences for heterogeneity both in meta-analysis of either experimental ( $Q=885.36$ ,  $p=0.001$ ,  $I^2=97.3\%$ ) or regular smoking ( $Q=369.22$ ,  $p=0.001$ ,  $I^2=96.5\%$ ). To deal with heterogeneity the random-effect model was used. The pooled prevalence estimation for experimental smoking was 27 % (95% CI; CI: 24- 30 %) and for regular smoking was 7 % (5 -9 %). The highest prevalence of regular smoking was reported from Gilan province in 2004 (15 %, 13 -17 %). The lowest prevalence was reported from Kerman in 2001 (2 %, 1 to 4 %). The highest and lowest prevalence of experimental smoking was reported from Qazvin province in 2001 (43 %, 37 - 48 %) and Kerman in 2001 (8 %, 5 to 11 %) respectively. There was no evidence for significant publication bias for experimental smoking in adolescents (Egger's test: 3.92, 95% CI: –

3.32 to 11.18;  $p=0.27$ ). But, there was a significant publication bias for regular smoking (Egger's test: 9.41; 95% CI, 1.01 to 17.81;  $p=0.03$ ).

#### 4. DISCUSSION

The prevalence of experimental and regular smoking in Zanjan city was 23.4 and 10.8 % respectively. These findings were higher than previous reports from Tabriz (northwest of Iran), (Mohammadpoor Asl, et al., 2007) and Shiraz (south of Iran) (Alireza Ayatollahi, et al., 2005). The meta-analysis showed that 27% and 7% of Iranian high school student were experimental or regular smokers respectively. This finding is in agreement with the Iran Global Youth Tobacco Survey (GYTS, 2007), which reported a figure of 23.7% and 5.1 % for experimental and regular smoking in 13 – 15 year old boys (Azaripour, 2007). Compared to international reports, the prevalence of adolescent smoking was considerably small. The prevalence of regular smoking in the Czech Republic (Baska, et al., 2006), Russian Federation (Baska, et al., 2009), and UK (Delpisheh, et al., 2007) have reported to be from 12% to 34%.

Despite of legislative restrictions of sale tobacco products to adolescents in Iran, majority of them (78.2 %) revealed easy access to cigarettes which means that tobacco control legislation and constraint without intersectional collaboration has only a little impact on cigarette accessibility.

The ordinal logistic regression model showed that increased chance of smoking could be expected by increasing age. This finding is consisted with other studies (Tyas & Pederson, 1998) (Mayhew, Flay, & Mott, 2000).

In the present study, using Tramadol and Ritalin medications were considered as a risk factor for adolescent smoking as these medications are strictly controlled in Iran and only specialists in some specific diseases can prescribe them for patients. Some adolescents are using these drugs without formal prescription. Overall, 36.0 % and 4.8 % of students had reported saw Tramadol and Ritalin respectively. The regression model also supported the hypothesis that using Tramadol has a significant relationship with



adolescent smoking. A possible explanation for this might be the fact that Tramadol have stimulus effect in human (Duke, Bigelow, Lanier, & Strain, 2011) and simultaneous using of Tramadol and cigarette results to increase in metabolism of Tramadol and therefore transient improvement in some psychological characteristics such as self-esteem, sense of kindness, and high energy. The MANOVA test supported the hypothesis that Tramadol and cigarette smoking have significant ordinal interaction on self-esteem ( $p < 0.001$ ). Hence, from a cross-sectional view, with abuse of Tramadol, tendency to cigarette smoking may be increased and this process can be known as a gateway for future regular smoking for current experimental adolescent smokers.

It was found that risk taking behavior (Alireza Ayatollahi, et al., 2005) (A Mohammadpoorasl, et al., 2011) (Kaplan, et al., 2001) (A. Mohammadpoorasl, et al., 2011) and smoker friends (Lakon & Valente, 2012) (Harakeh & Vollebergh, 2012) are significantly associated with adolescent smoking. A possible explanation for this might be due to adolescents' tendency of to have friends with similar smoking behavior (Mercken, Steglich, Sinclair, Holliday, & Moore, 2012). Recent studies revealed that “Passive (imitation) peer influence affected young adult smoking rather than active (pressure) peer influence” (Harakeh & Vollebergh, 2012).

It was found that relatives of experimental and regular smokers compared to never smokers have a mild reaction about lighting a cigarette in front of them. Other studies confirmed this result (Hanson, 1997) (Kaplan, et al., 2001) (Hrubá, Zaloudiková, & Matejová, 2010). But there is a little information that fears of relatives or family culture are effective.

There are some limitations in present study. 1. Due to nature of cross-sectional data, assessment of temporality between the related factors and cigarettes smoking was impossible. 2. To examine the relationship between Tramadol abuse and cigarette smoking, it was better to approve abuser people by biologic samples which it was not possible in the present study due to ethical issues. 3. The meta-analysis in Iran has some limitations especially in using standard search key words in national databases. To solve

this problem, all synonyms of search terms in both Persian and English languages were used.

In conclusion, the present study showed a moderately but concerning prevalence of experimental and regular adolescent smoking in Iran and introduced some novel associated risk factors. The effects of Tramadol abusing and its interaction with other psychological characteristic on adolescents' smoking habits should be investigated in future studies. In general, increasing prevalence of smoking in Iranian adolescents is a major concern for public health. Controlling risky behaviors and increasing health education are recommended.

## Acknowledgment

This research was performed with financial supports of Zanjan University of Medical Science, Zanjan, Iran, and Psychosocial Injuries Research Center, Ilam, Iran.

## References

- Aghaei A. (2001). Assessing the factors and attitudes to smoking in high school male students from the perspective of students, teachers and and parents in Isfahan. *Education Organization of Isfahan*.
- Ahmadi, J., & Hasani, M. (2003). Prevalence of substance use among Iranian high school students. *Addict Behav*, 28, 375-379.
- Ahmadi, J., & Ostovan, M. (2002). Substance use among Iranian male students. *International Journal of Drug Policy*, 13, 512-511.
- Alaei Khorayem R, Kadeyori P, Mohamadkhani Sh, Sarami GhR, & Alaei Khorayem S. (2011). The prevalence rate smoking, water pipe, alcohol, narcotics and stimulant among the high school students. *Journal of Research on Addiction*, 5, 114-199.
- Alireza Ayatollahi, S., Mohammadpoorasl, A., & Rajaeifard, A. (2005). Predicting the stages of smoking acquisition in the male students of Shiraz's high schools, 2003. *Nicotine Tob Res*, 7, 845-851.
- Aslinejad M, Alemi A, & Chamanzari H. (2004). Relationship between substance abuse with educational and family status of male high school students in Gonabad in the year 2003. In *Paper presented at the Third National Conference on Addiction*. Yazd: Iran.
- Azaripour, M. (2007). I.R.IRAN Global Tayth Tobacco Survey (GYTS) Reports 2007. In: WHO.
- Barikani A. (2008). High risk behaviors in adolescent students in Tehran. *Iran J Psychiatry Clin Psychol*, 14, 198-192.

- Baska, T., Sovinova, H., Nemeth, A., Przewozniak, K., Warren, C. W., & Kavcova, E. (2006). Findings from the Global Youth Tobacco Survey (GYTS) in Czech Republic, Hungary, Poland and Slovakia-- smoking initiation, prevalence of tobacco use and cessation. *Soz Praventivmed*, *51*, 110-116.
- Baska, T., Warren, C. W., Baskova, M., & Jones, N. R. (2009). Prevalence of youth cigarette smoking and selected social factors in 25 European countries: findings from the Global Youth Tobacco Survey. *Int J Public Health*, *54*, 439-445.
- Delpisheh, A., Attia, E., Drammond, S., & Brabin, B. J. (2006). Adolescent smoking in pregnancy and birth outcomes. *Eur J Public Health*, *16*, 168-172.
- Delpisheh, A., Kelly, Y., Rizwan, S., Attia, E., Drammond, S., & Brabin, B. J. (2007). Population attributable risk for adverse pregnancy outcomes related to smoking in adolescents and adults. *Public Health*, *121*, 861-868.
- Delpisheh, A., Kelly, Y., Rizwan, S., & Brabin, B. J. (2006). Socio-economic status, smoking during pregnancy and birth outcomes: an analysis of cross-sectional community studies in Liverpool (1993-2001). *J Child Health Care*, *10*, 140-148.
- Duke, A. N., Bigelow, G. E., Lanier, R. K., & Strain, E. C. (2011). Discriminative stimulus effects of tramadol in humans. *J Pharmacol Exp Ther*, *338*, 255-262.
- Epstein, D. H., Preston, K. L., & Jasinski, D. R. (2006). Abuse liability, behavioral pharmacology, and physical-dependence potential of opioids in humans and laboratory animals: lessons from tramadol. *Biol Psychol*, *73*, 90-99.
- Farajidana, H., Hassanian-Moghaddam, H., Zamani, N., & Sanaei-Zadeh, H. (2012). Tramadol-induced seizures and trauma. *Eur Rev Med Pharmacol Sci*, *16 Suppl 1*, 34-37.
- Habib E, Shirzad AS, Naseri-Kouzehgarani G, Hooman S, & Reza MM. (2012). The determinants of high school students smoking habits with special focus on teachers smoking in Iran: a population based study. *Pneumologia*, *61*, 33-28.
- Hamidzadeh Y, & Hosieni Ahog MM. (2001). Assess the prevalence smoking in the third year of high school boys students in 1998 Khalkhal. *J Ardabil Univ Med Sci*, *4*, 50-47.
- Hanson, M. J. (1997). The theory of planned behavior applied to cigarette smoking in African-American, Puerto Rican, and non-Hispanic white teenage females. *Nurs Res*, *46*, 155-162.
- Harakeh, Z., & Vollebergh, W. A. (2012). The impact of active and passive peer influence on young adult smoking: an experimental study. *Drug Alcohol Depend*, *121*, 220-223.
- Heydari, G., Sharifi, H., Hosseini, M., & Masjedi, M. R. (2007). Prevalence of smoking among high-school students of Tehran in 2003. *East Mediterr Health J*, *13*, 1017-1021.
- Hiemstra, M., Otten, R., & Engels, R. C. (2012). Smoking onset and the time-varying effects of self-efficacy, environmental smoking, and smoking-specific parenting by using discrete-time survival analysis. *J Behav Med*, *35*, 240-251.
- Hill, A. J., Boudreau, F., Amyot, E., Dery, D., & Godin, G. (1997). Predicting the stages of smoking acquisition according to the theory of planned behavior. *J Adolesc Health*, *21*, 107-115.
- Hruba, D., Zaloudikova, I., & Matejova, H. (2010). Similarities and differences between smoking and non-smoking ten-year-old children in primary schools. *Cent Eur J Public Health*, *18*, 19-24.
- Jafarzadegan T, & Zarehei F. (2001). The relationship between self-esteem and smoking among high school male students in Qazvin province. *Abstract of Research Report*.
- Kaplan, C. P., Napoles-Springer, A., Stewart, S. L., & Perez-Stable, E. J. (2001). Smoking acquisition among adolescents and young Latinas: the role of socioenvironmental and personal factors. *Addict Behav*, *26*, 531-550.
- Kelishadi, R., Ardalan, G., Gheiratmand, R., Majdzadeh, R., Delavari, A., Heshmat, R., Mokhtari, M. R., Razaghi, E. M., Motaghian, M., Ahangar-Nazari, I., Mahmood-Arabi, M. S., & Barekati, H. (2006). Smoking behavior and its influencing factors in a national-representative sample of Iranian adolescents: CASPIAN study. *Prev Med*, *42*, 423-426.

- Lakon, C. M., & Valente, T. W. (2012). Social integration in friendship networks: the synergy of network structure and peer influence in relation to cigarette smoking among high risk adolescents. *Soc Sci Med*, 74, 1407-1417.
- Marechal, C., Honorat, R., & Claudet, I. (2011). Serotonin syndrome induced by tramadol intoxication in an 8-month-old infant. *Pediatr Neurol*, 44, 72-74.
- Mayhew, K. P., Flay, B. R., & Mott, J. A. (2000). Stages in the development of adolescent smoking. *Drug Alcohol Depend*, 59 Suppl 1, S61-81.
- Mercken, L., Steglich, C., Sinclair, P., Holliday, J., & Moore, L. (2012). A longitudinal social network analysis of peer influence, peer selection, and smoking behavior among adolescents in British schools. *Health Psychol*.
- Miri M, Moasheri B, & Hanafi H. (2004). Prevalence and correlated factors of smoking among male students in Birjand and providing a suitable training model. In *Paper presented at the Third National Conference on Addiction*. Yazd: Iran.
- Mohammadpoor Asl, A., Fakhari, A., Rostami, F., & Pourafkary, N. (2007). Cigarette Smoking among Iranian Adolescents. *IJPBS*, 1, 30-35.
- Mohammadpoorasl, A., Fakhari, A., Shamsipour, M., Rostami, F., & Rashidian, H. (2011). Transitions between the stages of smoking in Iranian adolescents. *Prev Med*, 52, 136-138.
- Mohammadpoorasl, A., Nedjat, S., Fakhari, A., Yazdani, K., Rahimi Froushani, A., & Fotouhi, A. (2011). Smoking Stages in an Iranian Adolescent Population. In press.
- Mohtasham Amiri Z, Cirus Bakht S & NikRavesh Rad R (2008). Cigarette smoking in male high school students city of Rasht. *J Guilan Univ Med Sci*, 4, 100-107.
- Mojahed A, & Bakhshani Nm. (2004). Prevalence of smoking and drug abuse in students of Zahedan high schools. *J Mazandaran Univ Med Sci*, 6, 64-59.
- Mokhtari Habashi Gh, Bahrani N, & SobhRakhshankhah M. (2001). Prevalence of related factors and consequences of using addictive drugs among high school students in Urmia in the year 2001. *Urmia Med J*, 4, 40-51.
- Najafi K, Fekri F, & etal. (2007). Survey of the Prevalence of Substance Use Among High School Students In Guilan2005-2006. *J Guilan Univ Med Sci*, 16, 79-67.
- Najafi, K., Khalkhali, S., Nazifi, F., Farrahi, H., Faghripour, M., & Avakh, F. (2004). Prevalence of drug use in high school students in Rasht in 2003-04. In *Third National conference on Addiction*. Yazd: Iran, 2004.
- Namakin K, Sharifzadeh Gh, & Miri MR. (2008). Prevalence of cigarette smoking and evaluation of attitude and knowledge in its high school boys in Birjand, 2005. *J Birjand Univ Med Sci*, 15, 66-71.
- Rahmaniyan KA, Jafarzadeh A, & Khaloei A. (2010). Determining factors of smoking in high school students of Jahrom city. *Payavard Salamat*, 4, 96-88.
- Ramzankhani A, Sarbandi Zaboli F, Zargi A, Heydari GhR, & Masjedi MR. (2010). Smoking pattern in adolescent students of Tehran city. *Pejouhandeh*, 15, 122-115.
- Schaefer, D. R., Haas, S. A., & Bishop, N. J. (2012). A Dynamic Model of US Adolescents' Smoking and Friendship Networks. *Am J Public Health*, 102, e12-18.
- Shariat Zadeh M, & Sadeghi A. (2001). Epidemiological study of cigarette smoking in Urmia high school boys. *Med J Tabriz Univ Med Sci*, 34, 32-27.
- Tyas, S. L., & Pederson, L. L. (1998). Psychosocial factors related to adolescent smoking: a critical review of the literature. *Tob Control*, 7, 409-420.
- Wang, M. P., Ho, S. Y., Lo, W. S., & Lam, T. H. (2012). Smoking Family, Secondhand Smoke Exposure at Home, and Quitting in Adolescent Smokers. *Nicotine Tob Res*.

- Ziaaddini H, Kheradmand A, Nakhaei N, & Taherzadeh H. (2008). The prevalence of smoking and related factors in Kerman high school students. *The Quarterly Journal of Fundamentals of Mental Health*, 10, 245-239.
- Ziaei P, H. Z. N., Vameghi R, Dolatabadi S,. (2001). A study on prevalence of cigarette smoking and the age of first smoking in senior high school students in Tehran, 1998-99. *Hakim*, 4, 84-78.

**Table 1** Demographic characteristics and smoking status

Characteristics	Never smoker n (%)	Experimenter <sup>a</sup> n (%)	Regular smoker <sup>b</sup> n (%)	Total n (%)	p-value
Age					
≤15 years	61(81.3)	10(13.3)	4(5.3)	75(7.2)	P<0.001
16 years	159(67.1)	61(25.7)	17(7.2)	237(22.9)	
17 years	227(75.7)	51(17.0)	22(7.3)	300(29.0)	
18 years	141(58.0)	120(28.4)	69(28.4)	243(23.5)	
19 years	72(52.6)	40(29.2)	25(18.2)	137(13.2)	
≥20	21(48.8)	11(25.6)	11(25.6)	43(4.2)	
Total	681(65.8)	242(23.4)	112(10.8)	1035(100)	
Living with parents					
Yes	614(66.3)	211(22.8)	101(10.9)	926(92.2)	P<0.39
No	48(61.5)	23(29.5)	7(9.0)	78(7.8)	
Number of smoker friends					
0	464(77.5)	111(18.5)	24(4.0)	599(60.1)	P<0.001
≥1	190(47.7)	123(30.9)	85(21.4)	398(39.9)	
General risk-taking behavior					
Yes	454(61.9)	194(26.5)	85(11.6)	733(72.2)	P<0.001
No	216(76.6)	42(14.9)	24(8.5)	282(27.8)	
Socioeconomic status					
High class	158(67.2)	54(23.0)	23(9.8)	235(25.1)	P<0.87
Middle class	302(65.2)	110(23.8)	51(11.0)	463(49.4)	
Low class	157(65.7)	52(21.8)	30(12.6)	239(25.5)	
Death of family member					
Yes	57(58.8)	23(23.7)	17(17.5)	97(9.9)	P<0.06
No	593(67.2)	202(22.9)	88(10.0)	883(90.1)	
Membership in a sport team					
Yes	294(65.5)	94(20.9)	61(13.6)	449(45.6)	P<0.01
No	356(66.4)	135(25.2)	45(8.4)	536(54.4)	
Ever seen Tramadol					
Yes	181(48.5)	119(31.9)	73(19.6)	373(36.0)	P<0.001
No	500(75.5)	123(18.6)	39(5.9)	662(64.0)	
Ever seen Ritalin					
Yes	26(52.0)	10(20.0)	14(28.0)	50(4.8)	P<0.001
No	655(66.5)	232(23.6)	98(9.9)	985(95.2)	
Ever seen Ecstasy					
Yes	49(44.5)	32(29.1)	29(26.4)	110(10.6)	P<0.001
No	631(68.3)	210(22.7)	83(9.0)	924(89.4)	
Father norm					
Middle reaction	23(38.3)	19(31.7)	18(30.0)	60(6.1)	P<0.001
Severe reaction	622(67.7)	210(22.9)	87(9.5)	919(93.9)	
Mother norm					
Middle reaction	23(46.0)	11(22.0)	16(32.0)	50(5.0)	P<0.001
Severe reaction	637(67.1)	223(23.5)	90(9.5)	950(95.0)	
Brother norm					
Middle reaction	51(39.8)	44(34.4)	33(25.8)	128(18.6)	P<0.001
Severe reaction	401(71.7)	114(20.4)	44(7.9)	559(81.4)	
Sister norm					
Middle reaction	49(43.4)	38(33.6)	26(23.0)	113(15.7)	P<0.001
Severe reaction	428(70.6)	130(21.5)	48(7.9)	606(84.3)	
friend norm					
Middle reaction	279(55.4)	143(28.4)	82(16.3)	504(50.8)	P<0.001
Severe reaction	372(76.2)	90(18.4)	26(5.3)	488(49.2)	

a. Experimenters: respondents who indicated having tried or experimented with cigarette smoking, even a few puffs, but have smoked less than 100 cigarettes.

b. Regular smokers: respondents who indicated smoking 100 cigarettes or more in their lifetime irrespective of current smoking status.

**Table 2** Odds ratios (OR) and 95% confidence intervals (95% CI) of smoking of adolescents for related factors, based on logistic regression models.

Variables	Univariate model		Multivariate model	
	OR(95% CI)	P-value	OR(95% CI)	P-value
Ever use of Tramadol (yes)	3.27(2.50-4.28)	0.001	2.19(1.54-3.11)	0.001
Age (year)	1.30(1.18-1.44)	0.001	1.20(1.05-1.37)	0.005
Self-esteem (decrease)	1.08(1.06-1.11)	0.001	1.07(1.03-1.11)	0.001
Having general risk taking behaviors (yes)	3.46(2.55-4.70)	0.001	1.83(1.25-2.68)	0.002
Attitude toward smoking (increase)	1.22(1.17-1.27)	0.001	1.15(1.09-1.21)	0.001
Positive thinking about smoking (increase)	1.23(1.18-1.29)	0.001	1.07(1.01-1.14)	0.01
Having a smoker friend (yes)	3.76(2.85-4.95)	0.001	1.94(1.36-2.77)	0.001

**Author disclosure****Statement 1: Role of Funding Sources**

This research is performed with support of Zanjan University of Medical Science (ZUMS), Iran and Ilam University of Medical Science (IUMS), Iran . ZUMS and IUMS had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

**Statement 2: Contributors**

Milad Nazarzadeh, Dr.Ali delpishe, Ashgar Mohammadpoorasl, Abolfazl Bahrami, Zeinab Bidel and Erfan Ayubi designed the study and wrote the protocol. Zeinab Bidel, Erfan Ayubi and Milad Nazarzadeh conducted literature searches and provided summaries of previous research studies and study selection (especially for meta-analysis). Milad Nazarzadeh conducted the statistical analysis. Milad Nazarzadeh, Ali Delpisheh , Zeinab Bidel and Fateme Jafari wrote the first draft of the manuscript. Farhad Taromian and Ashgar Mohammadpoorasl was our consultant in addiction and psychologic topics. Abolfazl Bahrami manages data collection procedure and Erfan Ayubi data entry procedure. All authors contributed to critical appraisal of article and have approved the final manuscript.

**Statement 3: Conflict of Interest**

All other authors declare that they have no conflicts of interest.

**Statement 4: Acknowledgements**

The authors wish to thank Dr. Siamak Sabour, Shahid Beheshti University of Medical Sciences, Iran, for his valuable critical appraisal on this article; Dr. Masoud Vakili for his supports; Hadi Vaezi, Mehdi Bagheri, Reza Najafi and Asaad Kohi for their dedicated efforts in data collection procedures, Mohsen Rahimi for data entry procedure. We also thank all of the students and teachers of Zanjan city high schools for their valuable collaboration with this study.



#### Highlights

- Meta-analysis of smoking prevalence includes all Iranian high school students.
- Cross-sectional examines of psychological, pharmaceutical and social related factors with smoking.
- Tramadol can be a noble gateway for smoking.
- Adolescents' smoking is much more strongly associated with their peer's smoking.
- Relatives of experimenter and regular smoker have mild reaction about lighting a cigarette in front of them.